

PATENT APPLICATION

Our Docket No. 20020113.ORI

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Re App : Fionula M. Brennan et al : March 3, 2008
S.N. : 10/088,801 : Group Art Unit 1644
Filed : September 18, 2002 : Examiner Zachary Skelding
For : THERAPEUTIC METHODS AND COMPOUNDS

LETTER

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

Suggested new claims indicated as allowable by the Examiner
are attached for approval.

Respectfully submitted,

NIKOLAI & MERSEREAU, P.A.

C. G. Mersereau
Registration No. 26205
900 Second Avenue South
Suite 820
Minneapolis, MN 55402
Telephone: (612) 339-7461

132(new). A method of identifying a compound that selectively inhibits cytokine-stimulated T cell (T_{ck} cell) induced production of $TNF\alpha$ by monocytes at least two-fold greater than the compound inhibits T cell antigen receptor-stimulated T cell (T_{tor} cell) induced production of $TNF\alpha$ a by monocytes

wherein said method comprises the following steps:

- (a) providing a population of T_{ck} cells wherein said T_{ck} cells have been produced by incubating normal human peripheral blood T cells with cytokines selected from the group consisting of:
 - (i) IL-15;
 - (ii) IL-6, $TNF\alpha$ and IL-2; and
 - (iii) IL-6, $TNF\alpha$ and IL-15;
- (b) providing a population of T_{tor} cells wherein said T_{tor} cells have been produced by incubating normal human peripheral blood T cells with one or more anti-T cell antigen receptor antibodies which activate T cell antigen receptors;
- (c) incubating separate cultures of T_{ck} cells and T_{tor} cells with a compound to be tested;
- (d) resuspending the cultures of T_{ck} cells and T_{tor} cells in the absence of the test compound;
- (e) fixing the cultures of T_{ck} cells and T_{tor} cells;

(f) co-culturing equal portions of each of said fixed T_{ck} cells and T_{tcr} cells with monocytes to allow stimulation of the monocytes; and

(g) assaying for TNF α production by said monocytes, wherein the method identifies a compound that selectively inhibits T_{ck} cell induced production of TNF α by monocytes at least two-fold greater than the compound inhibits T_{tcr} cell induced production of TNF α by monocytes.

133(new). A method according to claim 132 wherein said compound is an antibody having specificity for T_{ck} cells.

134(new). A method according to claim 132 wherein said compound has efficacy in the treatment of rheumatoid arthritis.

135(new). A method according to claim 132 wherein said anti-T cell antigen receptor antibody is an anti-CD3 antibody.